

SINGLE VALUED CONJUGATES AND HOLOMORPHIC EXTENDIBILITY

Josip Globevnik

ABSTRACT Let D be a bounded domain in the complex plane whose boundary consists of $m \geq 2$ pairwise disjoint simple closed curves and let $A(bD)$ be the algebra of all continuous functions on bD which extend holomorphically through D . We show that a continuous function Φ on bD belongs to $A(bD)$ if for each $g \in A(bD)$ the harmonic extension of $\Re(g\Phi)$ to D has a single valued conjugate.

Let D be a bounded domain in the complex plane whose boundary consists of m pairwise disjoint simple closed curves. A continuous real function φ on bD has a unique continuous extension $\mathcal{H}(\varphi)$ to $\overline{D}$ which is harmonic on D . The conjugate $\mathcal{H}(\varphi)^*$ of $\mathcal{H}(\varphi)$ is a harmonic function on D which, in general, is multiple valued. It is determined up to an additive constant. Let $A(bD)$ be the algebra of all continuous functions on bD which extend holomorphically through D .

If $\Phi \in A(bD)$ then the harmonic extension of $\Re\Phi$, the real part of Φ , has the single valued conjugate $\Im\tilde{\Phi}$ where $\tilde{\Phi}$ is the holomorphic extension of Φ through D . Thus, for any $g \in A(bD)$, the harmonic extension of $\Re(g\Phi)$ has a single valued conjugate. In the present note we show that in the case when D is multiply connected, that is, when $m \geq 2$ this property characterizes the functions from $A(bD)$.

PROPOSITION *Let D be a bounded domain in the plane whose boundary consists of $m \geq 2$ pairwise disjoint simple closed curves. A continuous function Φ on bD extends holomorphically through D if and only if for each $g \in A(bD)$ the conjugate of the harmonic extension of $\Re(g\Phi)$ is single valued.*

In other words, Φ extends holomorphically through D if and only if for each $g \in A(bD)$ the harmonic extension of $\Re(g\Phi)$ to D is the real part of a single valued holomorphic function on D .

There is no such proposition in the case when D is simply connected as in this case every harmonic function on D has a single valued conjugate.

Every bounded domain $D \subset \mathbb{C}$ whose boundary consists of finitely many pairwise disjoint simple closed curves is biholomorphically equivalent to a domain D' bounded by finitely many pairwise disjoint circles [4]. Moreover, every biholomorphic map $\Phi: D \rightarrow D'$ extends to a homeomorphism $\tilde{\Phi}: \overline{D} \rightarrow \overline{D}'$ (see the proof in [2, pp. 46-49] which works also for multiply connected domains, bounded by finitely many pairwise disjoint simple closed curves). Thus, with no loss of generality assume that D is bounded by finitely many

pairwise disjoint circles $\gamma_1, \gamma_2, \dots, \gamma_m$, $m \geq 2$ where γ_m bounds the unbounded component of $\mathbb{C} \setminus \overline{D}$. For each j , $1 \leq j \leq m$, let h_j be the continuous function on $\overline{D}$, harmonic on D , such that $h_j|_{\gamma_j} \equiv 1$, $h_j|_{\gamma_k} \equiv 0$ ($1 \leq k \leq m$, $k \neq j$). These functions extend harmonically across bD . For each j , $1 \leq j \leq m$ let $\gamma'_j \subset D$ be a circle that has the same center as γ_j and that is a slight perturbation of γ_j . If u is a real harmonic function on a neighbourhood of $\overline{D}$ then the period of u^* along γ'_j is the same as the period of u^* along γ_j , that is, equal to $\int_{bD} \varphi \frac{\partial h_j}{\partial n} ds$ [3, p.80]. Since every real continuous function on $\overline{D}$ which is harmonic on D is the uniform limit of a sequence u_n such that each u_n is harmonic on a neighbourhood of $\overline{D}$ [3, p. 79] the preceding discussion implies that if φ is a real valued continuous function on bD then the period of $\mathcal{H}(\varphi)^*$ along γ'_j equals $\int_{bD} \varphi \frac{\partial h_j}{\partial n} ds$ so $\mathcal{H}(\varphi)^*$ is single valued if and only if

$$\int_{bD} \varphi \frac{\partial h_j}{\partial n} ds = 0 \quad (1 \leq j \leq m-1). \quad (1)$$

For each j , $1 \leq j \leq m$, $h_j + ih_j^*$ is a multiple valued holomorphic function in a neighbourhood of $\overline{D}$ whose complex derivative $W_j = (h_j + ih_j^*)'$ is a single valued holomorphic function in a neighbourhood of $\overline{D}$ without zeros on bD [3, p.81]. The Cauchy-Riemann equations imply that on bD

$$\frac{\partial h_j}{\partial n} ds = iW_j(\zeta)d\zeta \quad (1 \leq j \leq m)$$

[3, p.81]. Hence, by (1), $\mathcal{H}(\varphi)^*$ is single valued if and only if

$$\int_{bD} \varphi(\zeta)iW_j(\zeta)d\zeta = 0 \quad (1 \leq j \leq m-1). \quad (2)$$

The only if part of the proposition follows from the discussion preceding the proposition. To prove the if part, assume that Φ is a continuous function on bD such that for each $g \in A(bD)$ the conjugate of the harmonic extension of $\Re(g\Phi)$ is single valued.

By (2) it follows that for each $g \in A(bD)$ we have

$$\int_{bD} \Re[g(\zeta)\Phi(\zeta)]iW_j(\zeta)d\zeta = 0 \quad (1 \leq j \leq m-1)$$

which, since $iW_j(\zeta)d\zeta$ is real on bD , $1 \leq j \leq m-1$, implies that

$$\Re \left[\int_{bD} g(\zeta)\Phi(\zeta)iW_j(\zeta)d\zeta \right] = 0 \quad (1 \leq j \leq m-1).$$

It follows that for each $z \in \mathbb{C} \setminus \overline{D}$ we have

$$\int_{bD} \frac{1}{\zeta - z} \Phi(\zeta)W_j(\zeta)d\zeta = 0 \quad (1 \leq j \leq m-1)$$

which implies that for each j , $1 \leq j \leq m-1$, the function $\zeta \mapsto \Phi(\zeta)W_j(\zeta)$ extends holomorphically through D . So, for each j , $1 \leq j \leq m-1$, there is a function $H_j \in A(bD)$ such that

$$\Phi(\zeta) = \frac{H_j(\zeta)}{W_j(\zeta)} \quad (\zeta \in bD, 1 \leq j \leq m-1) \quad (3)$$

which means that for each j , $1 \leq j \leq m-1$, the function Φ extends through D as a meromorphic function with possible poles at the zeros of the function W_j . If we show that W_j , $1 \leq j \leq m-1$, have no common zero on D then (3) will imply that Φ extends holomorphically through D .

Fix $a \in D$ so close to bD that all the zeros $a_1, a_2, \dots, a_{m-1}$ of the Szegő kernel $z \mapsto S(z, a)$ are simple [1, p.106]. If $w \in D$ then the Garabedian kernel [1, p.24] is a meromorphic function on D with a single simple pole at w [1, p.25] which has no zero on $D \setminus \{w\}$ [1, p.49]. We now use the fact that the complex linear span of $\{W_1, W_2, \dots, W_{m-1}\}$ coincides with the complex linear span of $\{L(z, a_j)S(z, a) : 1 \leq j \leq m-1\}$ [1, p.80]. The preceding discussion implies that for each j , $1 \leq j \leq m-1$, the function $L(z, a_j)S(z, a)$ has zeros precisely at a_k , $1 \leq k \leq m-1$, $k \neq j$, which implies that the functions $L(z, a_j)S(z, a)$, $1 \leq j \leq m-1$ have no common zero on D . Suppose that $W_j(b) = 0$ ($1 \leq j \leq m-1$) for some $b \in D$. Then every function in the complex linear span of $\{W_1, W_2, \dots, W_{m-1}\}$ vanishes at b . In particular, all functions $L(z, a_j)S(z, a)$, $1 \leq j \leq m-1$, vanish at b , a contradiction. This shows that the functions W_j , $1 \leq j \leq m-1$, have no common zero on D . The proof of the proposition is complete.

We conclude with an example. Let $0 < R < 1$ and let $D = \{\zeta : R < |\zeta| < 1\}$. Let $\Phi = \varphi|_{bD}$ where $\varphi(z) = \bar{z}$. Let $g(z) = z^n$, $n \in \mathbb{Z}$, $n \neq 1$. Then $\Re(g\Phi)(e^{i\theta}) = \cos((n-1)\theta)$, $\Re(g\Phi)(Re^{i\theta}) = R^{n+1} \cos((n-1)\theta)$ ($0 \leq \theta < 2\pi$). The function

$$u(z) = \Re \left[\frac{R^{n+1} - R^{n-1}}{R^{n-1} - R^{1-n}} (z^{n-1} - z^{1-n}) + z^{n-1} \right]$$

is harmonic on $\mathbb{C} \setminus \{0\}$ and satisfies $u(e^{i\theta}) = \cos((n-1)\theta)$, $u(Re^{i\theta}) = R^{n+1} \cos((n-1)\theta)$ ($0 \leq \theta < 2\pi$) so u provides a harmonic extension of the function $\zeta \mapsto \Re[\zeta^n \Phi(\zeta)]$ ($\zeta \in bD$) to $\mathbb{C} \setminus \{0\}$. Clearly

$$u^*(z) = \Im \left[\frac{R^{n+1} - R^{n-1}}{R^{n-1} - R^{1-n}} (z^{n-1} - z^{1-n}) + z^{n-1} \right]$$

is single valued. Thus, for every $n \in \mathbb{Z}$, $n \neq 1$, the conjugate of the harmonic extension of the function $\zeta \mapsto \Re[\zeta^n \Phi(\zeta)]$ ($\zeta \in bD$) is single valued on D , yet Φ does not extend holomorphically through D .

This work was supported in part by the Ministry of Higher Education, Science and Technology of Slovenia through the research program Analysis and Geometry, Contract No. P1-0291.

REFERENCES

1. S. Bell: *The Cauchy Transform, Potential Theory and Conformal Mapping*. CRC Press, Boca Raton - Ann Arbor - London - Tokyo 1992
2. E. F. Collingwood and A. J. Lohwater: *The Theory of Cluster Sets*. Cambridge University Press, Cambridge 1966
3. S. D. Fisher: *Function Theory on Planar Domains*. John Wiley & Sons, New York 1983
4. G. M. Goluzin: *Geometrische Funktionentheorie*. VEB Deutscher Verlag der Wissenschaften, Berlin 1957

Institute of Mathematics, Physics and Mechanics
University of Ljubljana
Ljubljana, Slovenia
josip.globevnik@fmf.uni-lj.si